

FORM PTO-1390
(REV. 11-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

UBCP017

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/763298

INTERNATIONAL APPLICATION NO.

PCT/CA99/00813

INTERNATIONAL FILING DATE

September 3, 1999

PRIORITY DATE CLAIMED

September 3, 1998

TITLE OF INVENTION

Method For The Identification and Speciation of Bacteria of . . .

APPLICANT(S) FOR DO/EO/US

Mahenthiralingam

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

EL556131500US

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 20 below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
14. ☐ A SECOND or SUBSEQUENT preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
18. ☒ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☒ Other items or information:

Copy of IPER

Notification of Receipt of Demand

Notice Informing Applicant of Communication of Int. Application

INTERNATIONAL APPLICATION NO.

PCT/CA99/00813

ATTORNEY'S DOCKET NUMBER

UBCP017

21. ☒ The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO..... **\$1000.00**

International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO **\$860.00**

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO	\$710.00
---	-----------------

International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4)	\$690.00
--	----------

International preliminary examination fee (37 CFR 1.482) paid to USPTO
and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT =

CALCULATIONS PTO USE ONLY

\$ 860.00

Surcharge of **\$130.00** for furnishing the oath or declaration later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492(e)).

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$
--------	--------------	--------------	------	----

Total claims	$20 - 20 =$	0	x \$18.00	\$
--------------	-------------	---	-----------	----

Independent claims	$4 - 3 =$	1	x \$80.00	\$
--------------------	-----------	---	-----------	----

MULTIPLE DEPENDENT CLAIM(S) (if applicable)	+ \$270.00	\$
---	------------	----

TOTAL OF ABOVE CALCULATIONS	=	\$
------------------------------------	----------	-----------

<input checked="" type="checkbox"/>	Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.	- +	\$
-------------------------------------	---	--------	----

SUBTOTAL	=	\$
-----------------	---	----

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492(f)).

TOTAL NATIONAL FEE		=	\$
---------------------------	--	----------	-----------

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property	+	\$
---	---	----

TOTAL FEES ENCLOSED	=	\$
----------------------------	---	----

Amount to be refunded:

charged:

- a. ☒ A check in the amount of \$ 510.00 to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees.
A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 15-0610. A duplicate copy of this sheet is enclosed.
- d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO



PATENT TRADEMARK OFFICE

SIGNATURE

Marina T. Larson

NAME _____

PTO Reg. # 32,038

REGISTRATION NUMBER

UBCP017

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

U.S. Application No. To be assigned
Application of: Mahenthiralingam
International Application No. PCT/CA99/00813
International Filing Date: September 3, 1999
Priority Date Claimed: September 3, 1998
For: Method For The Identification and Speciation of Bacteria of the
BURKHOLDERIA CEPACIA Complex
Attorney Docket No. UBCP017

PRELIMINARY AMENDMENT

Asst. Commissioner for Patents
Washington, D.C. 20231

Sir:

Preliminary to the examination of the US national phase of PCT/CA99/00813 which is filed herewith, please make the following amendments to the claims, as amended before the IPEA/EP:

In the claims:

In claim 4, line 1, delete "any of claims 1 to 3" and insert -- claim 1 --.

In claim 11, line 3, delete "any of claims 8 - 10" and insert -- claim 8 --.

In claim 15, line 3, delete "claims 13 or 14" and insert -- claim 13 --.

09/763298

Please add claims 18, 19 and 20 as follows:

18. A kit for speciation of bacteria of the *Burkholderia cepacia* complex, comprising, in packaged combination, a pair of polynucleotide primers in accordance with claim 9, and a discriminatory restriction endonuclease.

19. A kit for speciation of bacteria of the *Burkholderia cepacia* complex, comprising, in packaged combination, a pair of polynucleotide primers in accordance with claim 10, and a discriminatory restriction endonuclease.

20. A kit for speciation of bacteria of the *Burkholderia cepacia* complex, comprising, in packaged combination, a pair of genomovar-specific polynucleotide primers in accordance with claim 14 and a discriminatory restriction endonuclease.

Respectfully,



Marina T. Larson
PTO Reg. No. 32,038
Attorney for Applicant
(970) 468-6600

METHOD FOR THE IDENTIFICATION AND SPECIATION OF
BACTERIA OF THE *BURKHOLDERIA CEPACIA* COMPLEX

Field of the Invention

This application relates to a new method for the identification and speciation of bacteria of the *Burkholderia cepacia* complex, and to vaccines developed specific for certain bacteria characterized using this method.

Background of the Invention

The Gram negative bacterium *Burkholderia cepacia* has recently been shown to consist of five different genomovars or new species, and as a collective the bacteria have been called the *B. cepacia* complex. Two of these genomovars have been given new species names: *Burkholderia multivorans* (formerly genomovar II) and *Burkholderia vietnamiensis* (formerly genomovar V). These bacteria cause problematic infections in patients with cystic fibrosis (CF) and chronic granulomatous disease as well as often causing infection outbreaks among vulnerable hospitalized patients. In cystic fibrosis, clinical outcome and epidemiology of infection may vary depending on the type of species patients are colonized with. It has become critical to obtain a rapid and reproducible means of identifying the different species of the *B. cepacia* complex since the current genomovar classification is technically difficult, laborious, carried out by only one reference laboratory and not based on a single test. Vandamme et al., *Int'l. J. Systematic Bacteriol.* 1188-1200 (1997).

It is an object of the present invention to provide a rapid and reproducible method for identification and speciation of *B. cepacia* complex.

It is a further object of the invention to provide suitable reagents and kits for use in the method of the invention.

Using this test, or other established but more time consuming tests, the species of a bacteria of the *B. cepacia* complex which is responsible for a given infection may be determined. There remains, however, the challenge of provide effective therapy if the strain is determined to be an epidemic strain. It is therefore still a further object of the present invention to provide an answer to this challenge by providing a vaccine which

- 2 -

promotes a therapeutically beneficial immune response to epidemic strains of the *B. cepacia*.

Summary of The Invention

In establishing a methodology for identification and speciation of a target group of microorganisms based on nucleotide sequences, it is necessary to identify a gene or genes within the target group which on the one hand contain conserved regions which are common to all microorganisms in the group such that non-specific amplification can be performed, and on the other hand contain regions which are dissimilar in ways that are diagnostic of the species. In accordance with the invention, it has been determined that these characteristics are found in the *recA* gene of bacteria of the *Burkholderia cepacia* complex. Thus, in one aspect of the invention, identification and speciation of bacteria of the *Burkholderia cepacia* complex in a sample can be accomplished by a method comprising the steps of

- (a) obtaining nucleotide sequence information for the *recA* gene in bacteria of the *Burkholderia cepacia* complex found in the sample; and
- (b) comparing the nucleotide sequence information obtained for the *recA* gene in bacteria of the *Burkholderia cepacia* complex found in the sample with a standard library of nucleotide sequence information comprising standard nucleotide sequence information for at least three species of bacteria of the *Burkholderia cepacia* complex.

Preferably, the nucleotide sequence information is obtained by evaluation of restriction fragment length polymorphism (RFLP). Other techniques for obtaining sequence information can also be used, including base-by-base determination of the sequence of the region of interest, sequence-specific oligonucleotide hybridization probes, and ligation techniques.

The invention also provides universal primers which can be used for amplification of all known members of the *Burkholderia cepacia* complex, and genomovar-specific primers which can be used for selective amplification of the *recA* gene from bacteria of one genomovar.

Speciation of bacteria of the *Burkholderia cepacia* complex can have several important consequences. When an infection has been identified as being caused by an

- 3 -

epidemic strain of the *B. cepacia* complex, it is appropriate to provide a vaccine which can be used in treating and preventing infection. Thus, the present invention also provides a vaccine composition which is useful in providing a therapeutically beneficial immune response in warm-blooded animals, particularly mammals (including humans) infected with epidemic strains of bacteria of the *B. cepacia* complex belonging to genomovar III. The vaccine comprises a protein or peptide antigen, or an expressible polynucleotide encoding a protein or peptide antigen derived from the flagellin of such bacteria. When the vaccine composition is administered to a mammalian subject, a therapeutically beneficial immune response is stimulated, which assists in combating the *B. cepacia* infection.

Speciation can also be used as a basis for selection and/or isolation of industrially useful bacterial species. For example, it is known that a number of strains of *B. cepacia* can be used in as biocontrol strains in agricultural applications. It has been found using the speciation method of the invention that these strains frequently fall into one of two classes based on *recA* RFLP pattern, one of which is a class associated with epidemic strains and one of which is not. Thus, one can use the test methodology of the invention to select among biocontrol strains to reduce the likelihood of public health hazards.

Brief Description of The Drawings

Figs. 1 and 2 show phylogenetic trees for *B. cepacia* based on the *recA* sequence evaluation; and

Fig. 3 shows the RFLP patterns obtained for representative strains of *B. cepacia* complex.

Detailed Description of The Invention

DNA sequence variation in conserved bacterial genes, such as those encoding the ribosomal RNA (rRNA) molecules, has been widely used as a method for classification and speciation of bacteria. Speciation of *B. cepacia* complex based on the genes encoding the 16S rRNA gene has been examined, but there is insufficient DNA variation within the gene to enable it to distinguish among all the genomovars of *B. cepacia*. In accordance

- 4 -

with the present invention, it has now been determined that there is sufficient sequence variation for speciation of *B. cepacia* in the gene encoding the RecA protein (*recA*). Thus, the present invention provides a rapid molecular diagnostic test which speciates bacteria of the *B. cepacia* complex based on the detection of sequence variation within the gene encoding the RecA protein.

The sequence of the *recA* gene has been evaluated in various bacterial species and suggested as a possible basis for phylogenetic classification. Karlin et al., *J. Bacteriol.* 177: 6881-6893 (1995). One sequence of *B. cepacia* was considered by Karlin et al., but there was no discussion of speciation. A similar phylogenetic comparison was published by Eisen, J.A., *J. Mol. Evol.* 41: 1105-1123 (1995), but again no comparisons among members of the *B. cepacia* complex were made.

The sequence of *recA* genes originating from two *B. cepacia* strains have been published. Nakazawa et al., *Gene* 94: 83-88 (1990); van Waasbergen, et al., *Appl. Microbiol. Biotech.* 49: 59-65 (1998). However, beyond noting that the sequences are "highly similar" but not identical, no particular utility was suggested as a result of the differences. Surprisingly, however, even though *recA* sequences have been shown in other cases to provide results similar to 16S rRNA, sequence evaluation of the *recA* gene of sixteen strains representative of each genomovar showed that these sequences contain sufficient diversity for speciation of the *B. cepacia* complex, while the 16S rRNA does not.

The published sequences for the *recA* gene in two strains sequences of bacteria of the *B. cepacia* complex (Seq. ID Nos. 1 and 2) were used to determine two primer sequences suitable for non-specific PCR amplification of the *recA* gene. This single pair of non-degenerate primers, which have the sequences:

Forward Primer (BCR1)

TGACCGCCGAGAAGAGCAA

SEQ ID No. 3

Reverse Primer (BCR 2)

CTCTTCTTCGTCCATCGCCTC

SEQ ID No. 4

were subsequently used to amplify the *recA* gene from additional strains, and were found to be effective as amplification primers for the *recA* gene from all known members of the *B. cepacia* complex, including LMG 14191^T, the type strain for *Burkholderia pyrrocinia*.

- 5 -

The amplified genes from these other strains were sequenced (Seq. ID Nos. 5-19), and the sequences aligned with the two previously known *recA* gene sequences. While there are substantial similarities, no two strains were identical except the sequences for 70431 and ATCC 17616. The *recA* genes from additional strains were sequenced and have strain numbers and GenBank Accession numbers, respectively, as follows: HI-2308, AF143777; C5424, AF143781; C1394, AF143783; ATCC 25416, AF143786; C2822, AF143792; LMG 10929, AF143793; LMG 14191 (*B. pyrrocinia*), AF143794; Ral-3, AF143795; M54, AF143796; M36, AF143797; ATCC 29424, AF143798; ATCC 53617, AF143799; ATCC 49709, AF143800; ATCC 39277, AF143801; ATCC 53266, AF143802.

The primers BCR1 and BCR2 are located at the 5'-ends of the sense and anti-sense strands, such that substantially the entire gene is amplified. Persons skilled in the art will appreciate, however, that other primers could be developed based on the sequences provided, for example to amplify only a portion of the gene in which mutations of diagnostic significance are found. Such alternative primers would be targeted to conserved regions of the *recA* gene, and could be degenerate if necessary to obtain amplification of all species within the *B. cepacia* complex.

Substantial blocks of nucleotide sequence which are conserved among all of the *recA* sequences determined are clearly from the aligned sequences. These nucleotide sequence regions may be used to design alternative primers for alternative amplification of *recA* from all bacteria of the *B. cepacia* complex. For example, there is substantial nucleotide variation from position 100 to 600 in the *recA* gene. Thus, a universal primer pair which amplified only this region would permit development of smaller amplified fragments effective for speciation. This can be accomplished using the primers:

Forward Primer (BCRU1*)

TGCGGATGGGCGACGGCG

SEQ ID No. 20

Reverse Primer (BCRU2*)

CAGTTCTGTCGCTTGATCG

SEQ ID No. 21

to produce a 485 base pair PCR product spanning a region of considerable sequence variation, or

Forward Primer (BCRU1)

ATCATGCGGATGGGCGACG

SEQ ID No. 36

- 6 -

Reverse Primer (BCRU2)

CAGTTCTGTCGCTTGATCG

SEQ ID No. 37

to produce a 488 base pair PCR product spanning a region of considerable sequence variation.

The DNA sequence variation in the *recA* gene used to speciate the *B. cepacia* can be detected using restriction enzyme digestion, separation of the digested fragments followed by pattern matching of the resulting profile. Prior to the digestion, the *recA* gene is amplified to increase the relative abundance of this gene as compared to other nucleic acid polymers in the sample. The preferred amplification technique is PCR amplification, but other known amplification techniques may be used as well. The restriction fragment patterns contain sufficient variation due to natural variation in the 16 *recA* sequences to enable the sequence to speciate among all 5 genomovars of *B. cepacia*. The enzymes *HaeIII* and *AluI* were found to be suitably discriminatory, with *HaeIII* providing the highest degree of discrimination. Other enzymes which might be possess adequate discrimination can be determined either by experimental comparison of RFLP patterns, or by computer analysis of the sequences to identify restriction sites based on the known cleavage sites of a given enzyme.

A phylogenetic tree based on the alignment of the novel sequences reported here and the two already published is shown in Figure 1 and demonstrates that the approach of the invention will clearly distinguish between all five of the current genomovars and also newly defined groups within the *B. cepacia* complex. This includes two sub-groups designated as RG-A and RG-B within genomovar III. The process of speciation of *B. cepacia* based upon RFLP using the *recA* gene scheme has been rigorously tested. It is highly discriminatory, reproducible and can be done using standard polymerase chain reaction techniques followed by agarose gel-electrophoresis.

It will be appreciated by persons skilled in the art, however, that while RFLP is a highly suitable method for evaluating nucleic acid sequence variations when (as in this case) enzymes can be identified which discriminate between the sequence variants, it is not the only method for making such a determination. Thus, nucleotide sequence information used in the method of the invention can be obtained by any other technique, including sequencing through at least the relevant regions of the *recA* gene; and the use of sequence-

- 7 -

specific hybridization probes or ligation techniques adapted to identify sequence variations. The 485 base pair fragment generated using primers BRCU1 and BRCU2 is particularly suited for direct sequencing, since its length is amenable to complete processing in conventional commercial sequencing apparatus. The primers BRCU1 and/or BRCU2 may also be used as sequencing primers. Comparison of this 485 bp fragment of the *B. cepacia* *recA* gene using phylogenetic tree software and alignment software is sufficient to separate the species clusters of the *B. cepacia* complex in the same fashion as comparison of the entire gene sequence. It is, however, a more rapid approach than determination of the entire gene sequence, since only one PCR product is involved which is of an optimal size for sequencing. Genomovar-specific RFLP of this fragment can also be performed to provide speciation of *B. cepacia* complex.

A further alternative for obtaining nucleotide sequence information indicative of the genomovar to which a sample bacterium of the *B. cepacia* complex belongs is through the use of PCR primers which are genomovar-specific. Such primers are selected such that at least one primer in the pair hybridizes to a region of the *recA* gene which is not conserved, i.e., to a variable region, in such a manner that amplification only occurs if the sequence of the variable region is complementary to the primer. These specific PCR primers enable a single PCR test to be used for identification and direct detection of strains of each genomovar.

The following are non-limiting examples of genomovar-specific primers. The specificity of these primers is such that other genomovars, other than the targeted genomovar, do not produce amplification products under stringent PCR conditions.

(i) *B. multivorans* specific *recA* primers:

Forward primer (BCRBM1):

5'- CGG CGT CAA CGT GCC GGA T - 3'

SEQ ID No. 22

Reverse primer (BCRBM2):

5' - T CCA TCG CCT CGG CTT CGT - 3'

SEQ ID No. 23

PCR product expected from *B. multivorans* strains = 714 bp.

(ii) *B. vietnamiensis* specific *recA* primers:

Forward primer (BCRBV1):

- 8 -

5'- GGG CGA CGG CGA CGT GAA - 3'

SEQ ID No. 24

Reverse primer (BCRBV2):

5' - TCG GCC TTC GGC ACC AGT - 3'

SEQ ID No. 25

PCR product expected from *B. vietnamiensis* strains = 378 bp.**(iii) *B. cepacia* Genomovar IV specific *recA* primers:**

Forward primer (BCRG41):

5' - ACC GGC GAG CAG GCG CTT - 3'

SEQ ID No. 26

Reverse primer (BCRG42):

5' - ACG CCA TCG GGC ATG GCA - 3'

SEQ ID No. 27

PCR product expected from *B. cepacia* strains of genomovar IV = 647 bp.**(iv) *B. cepacia* genomovar III, RG-B *recA* specific primers.**

Forward primer:

5' - GCA AGT CAT CGC TGA GAA - 3'

SEQ ID No. 28

or

Forward primer:

5' - GCT GCA AGT CAT CGC TGA A - 3'

SEQ ID No. 38

Reverse primer):

5' - TAC GCC ATC GGG CAT GCT - 3'

SEQ ID No. 29

PCR product expected from strains of this new genomovar classification = 781 bp.

(v) *B. cepacia* genomovar I specific *recA* primers:

Forward primer (BCRG11):

5' - CAG GTC GTC TCC ACG GGT - 3'

SEQ ID No. 30

Reverse primer (BCRG12):

5' - CAC GCC GAT CTT CAT ACG A - 3'

SEQ ID No. 31

PCR product expected from strains of genomovar I = 492 bp.

(vi) *B. cepacia* genomovar III, RG-A specific *recA* primers:

Forward primer (BCRG31)

5' - GCT CGA CGT TCA ATA TGC C - 3'

SEQ ID No. 32

Reverse primer (BCRG32):

T00220" B62E260

- 9 -

5' - TCG AGA CGC ACC GAC GAG - 3'

SEQ ID No. 33

PCR product expected from *B. cepacia* strains of genomovar III = 378 bp.

Additional sequencing of the complete *recA* gene from *B. cepacia* complex strains M36, M54, Ral-3 and *B. pyrrocinia* LMG 14191^T or partial sequence analysis of PCR amplicons derived from strains ATCC 29464, ATCC 53617, ATCC 39277, ATCC 49709 and ATCC 53266 was performed. The phylogenies determined using partial sequencing of this type were identical to those determined using the full sequences (Fig. 2), however, two additional clusters, denominated as RG-C and RG-D were identified. Cluster RG-C was a novel group consisting of biocontrol strains Ral-3, ATCC 53266 and M54. Comparative alignment of the *recA* sequence from Ral-3 and M54 with all other complete *B. cepacia* sequences enabled the design of RG-C specific primers with the following sequences:

Forward Primer:

GTCGGGTAAAACACGTG

SEQ ID No. 39

Reverse Primer:

TCCGCAGCCGCACCTTCA

SEQ ID No. 40

B. cepacia biocontrol strains BC-B, BC-F and AMMD all tested positive with this RG-C primer set. Thus, these primers can be used in analytical schemes for the presence of such primers, and also could be used for screening isolates for biocontrol properties.

A second novel *recA* group, RG-D, was identified which includes *B. pyrrocinia* LMG 14191^T and ATCC 32977, a strain of *B. cepacia* which produces the antibiotic xylocladin. This group is also shown in Fig. 2.

In addition to a method for identification and speciation of bacteria of the *B. cepacia* complex, the invention also provides reagents and kits suitable for carrying out this method. The reagents are generally polynucleotide primers or probes which bind to the *recA* gene of one or more strains of bacteria of the *B. cepacia* complex. One subset of the reagents of the invention are non-specific primers, such as used in Example 4 below, which are complementary to conserved regions found identically in strains of bacteria of the *B. cepacia* complex for which the sequences are given. A second subset of reagents in

- 10 -

accordance with the invention are primers/probes which can be used to selectively amplify and/or detect one genomovar of bacteria of the *B. cepacia* complex. The reagents of the invention may have a detectable or capturable label, for example a radioactive or fluorescent label or biotin, incorporated therein to facilitate evaluation of nucleotide sequence information.

Either of these types of primers/probes may be packaged in a kit with suitable reagents. These reagents may include discriminatory restriction enzymes, which are capable of producing distinctive fragment patterns to permit speciation of a bacteria-containing sample, or reagents suitable for PCR, nucleic acid sequencing and the like.

Once the species of a sample bacterium of the *B. cepacia* complex is determined using the method of the invention, it may be desirable (particularly where the bacteria is a member of an epidemic strain) to be able to provide a therapeutic agent which is effective in treating or preventing infection. Thus, the present invention further provides a vaccine composition based upon the antigenic properties of the flagellin of epidemic strains of *B. cepacia* complex for use in treating infections caused by certain species of the *B. cepacia* complex.

The use of flagellins as an antigen for vaccine purposes has been proposed in a variety of instances because of their location on the outside of bacterial cells. In the case of *B. cepacia* complex, however, Hales et al., *J. Bacteriol.* 180: 1110-1118 (1998), have reported that the flagellin gene (*fliC*) is "highly variable" and suggest its utilization as a biomarker for epidemiological and phylogenetic studies of *Burkholderia cepacia*. Such variability is inconsistent with the normal requirements that a vaccine antigen be highly conserved, such that its will be generally effective against variants of the target species. Thus, it was quite surprising to find that the subset of *B. cepacia* complex which is most transmissible have highly conserved flagellin genes which is suitable for use as a vaccine.

A total of 30 strains of bacteria of the *B. cepacia* complex were classified using the speciation method of the invention into groups based on the sequence of the *recA* gene, and were in addition characterized with respect to the BCESM (*Burkholderia cepacia* Epidemic Strain Marker) and *cbIA* markers for highly transmissible strains of *B. cepacia*. As reflected in Table 1, a substantial portion of the genomovar III strains which were positive for one or both of these markers produced a

- 11 -

single RFLP pattern (Fig. 3, pattern G) after treatment with the restriction endonuclease *HaeIII*.

Exemplary sequences and a consensus sequence for the *B. cepacia* flagellin gene, which encodes the major subunit protein of the bacterial flagellum of *B. cepacia*, have been described in the literature by Hales et al. (supra). Using the same primers described by Hales, it has been determined that the flagellin genes of *B. cepacia* strains of *recA* type III-G (genomovar III, with *recA* RFLP pattern G) are highly conserved and do not vary considerably in DNA sequence. This indicates that the protein is also highly conserved in its structure and sequence, and thus is suitable for use as an antigen for development of vaccines against the most problematic strains in patients with cystic fibrosis (CF).

In contrast, *B. multivorans* strains of *recA* type F (genomovar II), which appear less problematic in patients with CF and do not generally spread among patients, have flagellin genes which are highly variable in sequence. These data suggest that with *B. multivorans* strains, a vaccine based on the flagellum may not protect against infection with all strains types as has been the case with the bacterium *Pseudomonas aeruginosa* in CF. Thus, the observation, that the flagellin gene is actually highly conserved in the most devastating epidemic *B. cepacia* strains infecting patients with CF is apparently unique to this subset of the species of the *B. cepacia* complex.

The observation that the flagellin gene is conserved among *B. cepacia* strains which are epidemic amongst patients with CF permits the development of a vaccine based on the encoded protein antigen. The vaccine can be prepared in a variety of ways. First, protein can be purified from bacterial strains representative of this group to obtain a purified antigen. Methods for purification of flagellin from bacteria are known in the art, and can be applied to recovery of purified flagellin from epidemic strains of *B. cepacia*. Purified antigen is then used as a vaccine, with or without an adjuvant. Vaccines of this type are generally administered by subcutaneous or intramuscular injection, although other routes of administration may also be suitable. Therapeutically effective levels and frequency of vaccine administration are determined by routine monitoring of antibody titers.

In addition to the use of purified flagellin isolated directly from bacteria, it will be appreciated that the same protein, or an immunologically effective portion thereof may

- 12 -

also be prepared using, for example, recombinant technology. Thus, for example, cDNA encoding flagellin or an immunologically effective portion thereof may be cloned into a host organism and expressed to produce flagellin antigen. Smaller antigenic peptides may also be made synthetically. As used herein, the term "derived from" refers to proteins or peptides which are either isolated directly from bacteria of the *B. cepacia* complex, or which have the same amino acid sequence but which are obtained synthetically or by expression in a host organism.

Vaccine compositions in accordance with the invention comprise the flagellin or flagellin-derived antigen, in a pharmaceutically acceptable carrier. In general such carriers will be aqueous media, and may include buffers, emulsifiers, or adjuvants to enhance the immune response. Antigenic proteins or peptides may also be provided in association with lipid-carriers, e.g., liposomes or other lipid particles, in the vaccine composition.

The composition of the invention may also include other materials which are more indirectly derived from flagellin, and which provide immunoprotective therapeutic benefits. For example, human antibodies specific to *B. cepacia* flagellin or immunologically effective portion thereof are considered to be "derived-from" flagellin in accordance with the invention. Such antibodies may be administered to individuals with *B. cepacia*, genomovar III infections, to help combat the progression of infection. Such therapy is particularly suitable in end-stage *B. cepacia* infection when antibiotics and anti-inflammatory therapy have failed.

Vaccination may also be carried out using DNA vaccines of the type described generally in US Patent No. 5,580,859, which is incorporated herein by reference. DNA vaccines comprise a sequence encoding the desired protein or peptide antigen, optionally in combination with a regulatory element to control expression of the antigen. The DNA vaccine, which may be naked or incorporated in a carrier such as a liposome, is administered by subcutaneous or intramuscular injection. A "gene gun" may be used for administration.

- 13 -

EXAMPLE 1

Restriction fragment length polymorphisms (RFLP) of the 16S rRNA gene amplified from strains from the *B. cepacia* complex were determined. The amplified fragment of the 16S rRNA gene was digested with the enzyme *Dde* I and the resulting products separated by agarose gel electrophoresis. Species specific patterns were found for *B. vietnamiensis* (genomovar V) and *B. multivorans* (genomovar II) strains, however sequence variation in the 16S rRNA gene was insufficient to distinguish among strains from genomovars I, III and IV.

EXAMPLE 2

The novel DNA sequences of 16 *recA* genes, from 16 strains of the genomovar representative panel shown in Table 1, have now been obtained by conventional DNA sequence analysis of portions of the *recA* gene amplified by PCR. DNA sequence analysis was performed in collaboration with Prof. Julian Davies at TerraGen Diversity Inc., Vancouver, BC, using an ABI 377 Nucleotide Sequencer. The 1040 bp *recA* gene was split into two ~520 bp PCR products to facilitate complete sequence analysis. The primer pairs used were as follows:

(i) 5' - portion of the *recA* gene:

Forward primer BCR1:

5' - TGA CCG CCG AGA AGA GCA A- 3'

SEQ ID No. 3

Reverse primer BCR4:

5' - GCG CAG CGC CTG CGA CAT - 3'

SEQ ID No. 34

These primers amplified a 527 bp product corresponding to the 5' half of the *recA* gene from all members of the *B. cepacia* complex tested. Nucleotide sequence from both strands of the amplified products was determined by direct nucleotide sequence analysis using primers BCR1 and BCR4 respectively.

(ii) 3' - portion of the *recA* gene:

Forward primer BCR 3:

5' - GTC GCA GGC GCT GCG CAA - 3'

SEQ ID No. 35

Reverse primer BCR 2:

5' - CTC TTC TTC GTC CAT CGC CTC - 3'

SEQ ID No. 4

- 14 -

These primers amplified a 529 bp product corresponding to the 3' half of the *recA* gene from all members of the *B. cepacia* complex tested (see Table 1). Nucleotide sequence from both strands of the amplified products was determined by direct nucleotide sequence analysis using primers BCR2 and BCR3 respectively.

Because the amplified products described above overlapped in the regions corresponding to primers BCR3 and BCR 4, the complete 1040 bp *recA* gene sequences were derived by joining of each half of nucleotide sequence data. The alignment of these sequences was carried out in collaboration with Dr. Yossef Av-Gay, Dept. of Medicine, Vancouver General Hospital, Vancouver, BC, using the software CLUSTAL which is available on the internet. The two published *recA* sequences, b-cepd90120_1 and b-cepu70431_1 were aligned with 16 novel sequences obtained using the speciation scheme. Sequence b-cepk56-2_1 is the *recA* gene of a *B. cepacia* strain from the same cable-pilus gene encoding lineage cited in PCT Patent Applications Nos. WO 97/01647 and WO97/07237. From such a sequence alignment, the sequence variation in the *recA* gene detected by the Hae III RFLP analysis is clearly visible. These sequences can also be used for the design of genomovar-specific amplification/sequencing primers, and genomovar-specific hybridization or ligation probes.

EXAMPLE 3

A *B. cepacia* complex phylogenetic tree based on the sequence alignment of the novel *recA* genes determined was formulated as shown in Fig. 1. The alteration of single bases in the nucleotide sequence of genes by natural mutation over time can be quantitated by computer programs to an evolutionary distant which separates strains and species. Phylogenetic analysis of the *recA* sequences which I have determined clearly demonstrates that sequence variation in this gene can separate all five genomovars of the *B. cepacia* complex, shown by the tree in the figure. Even more interesting is the novel finding of two distinct groups within the strains otherwise classified within genomovar III. Thus, *recA* sequences derived from strains CEP511, C1394 and PC 184, respectively, (Table 1) cluster as a separate group on a different branch of the tree from *recA* sequences derived from strains C6433, C4455m K56-2 and C5425. These two groups have been designated a RG-B and RG-A, respectively. The three strains of the RG-B are all

- 15 -

epidemic amongst patients with CF (Mahenthiralingam et al., *J. Clin. Microbiol.* 34: 2914-2920 (1996) and encode the BCESM (Mahenthiralingam et al., *J. Clin. Microbiol.* 35: 808-816 (1997). The separate classification of these three strains based on the *recA* suggest that they may constitute a new species/genomovar group within the *B. cepacia* complex.

EXAMPLE 4

To obtain nucleotide sequence information about the *recA* genes of additional strains of bacteria of the *B. cepacia* complex (Table 1), samples of each strain were amplified using the following primers:

Forward Primer (BCR1)

TGACCGCCGAGAAGAGCAA

SEQ ID No. 3

Reverse Primer (BCR 2)

CTCTTCTTCGTCCATCGCCTC

SEQ ID No. 4

using a standard polymerase chain reaction mixture of 25 microlitres in volume (described in Mahenthiralingam et al., *J. Clin. Microbiol.* 35: 808-816 (1997)) containing 1.5 mM MgCl₂ and 10-20 ng of *B. cepacia* DNA. Amplification was performed as follows: 30 cycles of 1 min. at 94°C, 1 min. at 56°C, and 2 min. at 72°C, follow by a final 6 min. cycle at 72°C. This resulted in the amplification of a 1 kb DNA band corresponding to the *recA* gene of the *B. cepacia* strain being tested.

Several restriction enzyme were screened for their ability to reveal DNA sequence variation in this amplified gene which would be suitable for speciation of *B. cepacia*. The enzymes *Hae* III and *Alu* I were found to be suitably discriminatory. The restriction fragments produced by the enzyme *Hae* III were separated by agarose gel-electrophoresis, and the detected restriction fragment length polymorphisms (RFLPs) demonstrated that genomovar specific RFLPs could be generated using this approach. Representative patterns are shown in Fig. 3. (Bv = *B. vietnamiensis*, or genomovar V; Gv I = genomovar I; Bm = *B. multivorans* or genomovar II; Gv III = genomovar III and Gv IV = genomovar IV). This same approach has been applied to a panel of strains which are representative of all five genomovars of *B. cepacia* and been found to be able to distinguish strains of each genomovar (Table 1). This technique has also been applied to additional strains, and been

- 16 -

found to be reproducible and highly discriminatory for speciation of strains from the *B. cepacia* complex.

EXAMPLE 5

To obtain nucleotide sequence information about the *recA* genes of 30 strains of bacteria of the *B. cepacia* complex (Table 1), samples of each strain were amplified using the following primers:

Forward Primer (BCR1, %G/C Tm =53.2°C)

TGACCGCCGAGAAGAGCAA

SEQ ID No. 3

Reverse Primer (BCR 2, (%G/C Tm =56.3°C)

CTCTTCTTCGTCCATCGCCTC

SEQ ID No. 4

using a standard polymerase chain reaction mixture of 25 microlitres in volume (described in Mahenthiralingam et al., *J. Clin. Microbiol.* 35: 808-816 (1996)) containing 1.5 mM MgCl₂ and 10-20 ng of *B. cepacia* DNA. Amplification was performed as follows: 30 cycles of 1 min. at 94°C, 1 min. at 56°C, and 2 min. at 72°C, follow by a final 6 min. cycle at 72°C. This resulted in the amplification of a 1 kb DNA band corresponding to the *recA* gene of the *B. cepacia* strain being tested (Fig. 2A).

Several restriction enzyme were screened for their ability to reveal DNA sequence variation in this amplified gene which would be suitable for speciation of *B. cepacia*. The enzymes *Hae* III and *Alu* I were found to be suitably discriminatory. The restriction fragments produced by the enzyme *Hae* III were separated by agarose gel-electrophoresis (Fig. 2B) and the detected restriction fragment length polymorphisms (RFLPs) demonstrated that genomovar specific RFLPs could be generated using this approach (Bv = *B. vietnamiensis*, or genomovar V; Gv I = genomovar I; Bm = *B. multivorans* or genomovar II; Gv III = genomovar III and Gv IV = genomovar IV).

EXAMPLE 6

Bacterial strains identified in Example 5 as belonging to RFLP group G were epidemic strains of genomovar III which encode the BCESM DNA. When the flagellin gene of these strains was amplified using the PCR primers as described in Hales et al., a 1 kb product corresponding to the flagellin gene was obtained. Sequence variation

09763258-022001

- 17 -

in amplified flagellin gene was again detected using the restriction enzyme *Hae* III. The RFLP of the flagellin genes was highly conserved demonstrating that both the gene and the encoded protein are very conserved in sequence. The conservation in sequence of the flagellin of these epidemic strains suggests that the flagellum is an ideal candidate upon which to develop a vaccine able to protect against *B. cepacia* infection with these problematic strain types. This novel observation on the conservation of the flagellin gene and protein of epidemic *B. cepacia* strain types is in contrast to the variation reported in Hales et al. and that which is observed for *B. multivorans* strains (genomovar II) which is described in Example 7.

EXAMPLE 7

Using the same scheme outlined in Example 6, the flagellin gene in *B. multivorans* (genomovar II) strains which colonized patients attending clinic in Vancouver was examined. These strains were of a single *recA* type, BM-F, demonstrating that they are a single species type, *B. multivorans*. When the flagellin gene of these strains was examined using PCR amplification followed by RFLP analysis with *Hae* III, the gene of each of the strains examined was highly variable. This suggest that the flagellum may not be as useful in protection against infection with these strain types. However, infection with these *B. multivorans* appears not to be as problematic as infection with the epidemic *recA* III-G/BCESM positive strains.

EXAMPLE 8

Thirteen isolates having useful biological properties were evaluated to determine the *recA* group to which the isolates belong. The results of these and other tests are summarized in Table 2. Two strains within the ATCC collection with interesting catabolic properties (ATCC 29424 and ATCC 53617) were classified as *B. vietnamensis* on the basis of both 16S rRNA ARDRA and *recA* analysis. Of the remaining 11 commercially useful strains, all possessed ARDRA polymorphisms characteristic of *B. cepacia* genomovars I, III(RG-A and RG-B) and IV. Strain ATCC 49709, a seed-treatment biocontrol strain, was classified as genomovar I by *recA* RFLP and specific *recA*-PCR. Strains M36, BC-1 and BC-2, biocontrol strains isolated from the rhizosphere

- 18 -

of maize crops, possessed *recA* RFLP types characteristics of genomovar III, group RG-B and all tested positive in RG-B specific PCR. The remaining seven strains all possessed novel *recA* RFLP types and were not reactive for any of the genomovar/ sub group specific PCR primers described above.

Information of the type developed in this assay can potentially be used to assess the safety of strains of *B. cepacia* complex for commercial applications, including agricultural applications. Prior to this invention, no distinct criteria were established for division of pathogenic from non-pathogenic bacteria of the *B. cepacia* complex. However, the biologically useful strains tested were found to belong to several *recA* phylogenetic groups. Some strains cluster within the genomovar III, *recA* subgroup RG-B, which is the same subgroup to which a number of well-characterized pathogenic strains belong. This includes strain M36, which encodes the BCESM and which has been withdrawn from sale by the manufacturer. On the other hand, a number of the strains also belong to other subgroups which are not associated with pathogenicity. Thus, *recA* analysis may provide a simple method for screening biocontrol strains and other potentially useful members of the *B. cepacia* complex to identify strains that are less likely to raise health and public safety issues in their use. Furthermore, a number of the biocontrol strains were found to occupy a novel *recA* phylogenetic subgroup, RG-C. The RG-C specific PCR primers, or other tests based on the sequence analysis performed on clinical isolates indicate that RG-C strains are not encountered in human infections, and appear specifically adapted to the plant rhizosphere, they may prove to be the best template upon which to continue to development of safe biocontrol strains.

0976359-022001

TABLE 1
Provisional list of strains in the *B. cepacia* complex experimental panel

ID No.	Strain Name:	Genomovar:	Source: ^a	RAPD ^b type:	BCESM: ^c	<i>cblA</i> :	Gene RFLP type: <i>recA/Hae</i> III	16S rRNA/ <i>DdeI</i>
1	PC259	<i>B. vietnamiensis</i> (V)	CF	8	-	-	A	1
2	LMG 16232	<i>B. vietnamiensis</i> (V)	CF	X	-	-	A	1
3	ATCC 39277	<i>B. vietnamiensis</i> (V)	ENV	X	-	-	C	2
4	LMG 10929	<i>B. vietnamiensis</i> (V)	ENV	X	-	-	B	1
5	C2822	<i>B. vietnamiensis</i> (V)	CF	X	-	-	B	1
8	FC0441	<i>B. vietnamiensis</i> (V)	CGD	61	-	-	A	1
6	ATCC 25416	I	ENV	39	-	-	D	2
7	CEP0509	I	CF	41	-	+	E	2a
9	ATCC 17759	I	ENV	X	-	-	E	2a
10	C5393	<i>B. multivorans</i> (II)	CF	3	-	-	F	3
11	C3430	<i>B. multivorans</i> (II)	CF	7	-	-	F	3
12	C5274	<i>B. multivorans</i> (II)	CF	12	-	-	F	3
13	C5568	<i>B. multivorans</i> (II)	CF	19	-	-	F	3
14	249-2	<i>B. multivorans</i> (II)	LAB	30	-	-	F	3
15	ATCC 17616	<i>B. multivorans</i> (II)	ENV	30	-	-	F	3
16	LMG 13010	<i>B. multivorans</i> (II)	CF	X	-	-	F	3
17	JTC	<i>B. multivorans</i> (II)	CGD	X	-	-	F	3
18	C1257	III	CF-e	1	+	-	G	2
19	BC7	III	CF-e	2	+	+	G	2
20	J2315	III	CF-e	2	+	+	G	2
21	C5424	III	CF-e	2	+	+	G	2
22	LMG 12615	III	CF-e	2	+	+	G	2
23	C6433	III	CF-e	4	+	-	G	2
24	C4455	III	CF-e	6	+	-	G	2
25	C1394	III	CF-e	13	+	-	H	2
26	PC184	III	CF-e	17	+	-	J	2
27	CEP0511	III	CF-e	40	+	-	I	2
28	LMG 14291	IV	CF	X	-	-	J	2
29	LMG 07000	IV	CLIN	16	-	-	J	2
30	LMG 14294	IV	CF	16	-	-	J	2

Footnotes:

- ^a Source of isolate; CF, cystic fibrosis infection; CF-e, epidemic amongst CF patients (31); ENV, environment; CGD, infection of a chronic granulomatous disease patient; LAB, laboratory derived and CLIN, non-CF clinical infection.
- ^b RAPD type derived from Mahenthiralingam *et al.* (30): X=unique strain RAPD fingerprint unmatched in our collection.
- ^c BCESM and *cblA* hybridization data adapted from Mahenthiralingam *et al.* (31)

Table 2
Analysis of *B. cepacia* complex strains with useful biological properties

Strain:	Source and useful property ^a :	BCESM:	16S rDNA RFLP (<i>Dde</i> I):	<i>recA</i> (<i>Hae</i> II) RFLP:	<i>recA</i> PCR group:	Reference/ Origin
ATCC 29424	Soil isolate, capable of phthalate utilization	+	1	B	BV	51
ATCC 53617	Waste water isolate, trichloroethylene degrader	-	1	A	BV	ATCC
ATCC39277	Cornfield soil isolate, antifungal agent	-	2	P	Novel	34
ATCC 49709	Grass seed isolate, biocontrol strain	-	2	D	G1	ATCC
M36	Corn rhizosphere isolate, biocontrol strain	+	2	I	RG-B	Stine Seed Co.
BC-1 ^b	Corn rhizosphere isolate, biocontrol strain	+	2	H	RG-B	USDA
BC-2 ^b	Corn rhizosphere isolate, biocontrol strain	+	2	I	RG-B	USDA
M54	Corn rhizosphere isolate, biocontrol strain	+	2	L	RG-C	Stine Seed Co.
Ral-3	Corn rhizosphere isolate, biocontrol strain	-	2	N	RG-C	Agrium Inc.
ATCC 53266	Corn rhizosphere isolate, biocontrol strain	-	2	L	RG-C	ATCC
BC-B	Corn rhizosphere isolate, biocontrol strain	-	2	L	RG-C	USDA
BC-F ^c	Corn rhizosphere isolate, biocontrol strain	-	2	N	RG-C	USDA
AMMD	Pea rhizosphere isolate, biocontrol strain	-	2	N	RG-C	23

Footnotes:

^a Biocontrol strains demonstrated protection of crops against a various of phytopathogens; these included either fungal infection or nematode infections.

^b Strain originally isolated by K. Prakash Hebbar, USDA.

^c Strain originally isolated by WeiLi Mao, USDA.

- 21 -

Claims

1. A method for identification and speciation of bacteria of the *Burkholderia cepacia* complex in a sample, comprising the steps of
 - (a) obtaining nucleotide sequence information for the *recA* gene in bacteria of the *Burkholderia cepacia* complex found in the sample; and
 - (b) comparing the nucleotide sequence information obtained for the *recA* gene in bacteria of the *Burkholderia cepacia* complex found in the sample with a standard library of nucleotide sequence information comprising standard nucleotide sequence information for at least three species of bacteria of the *Burkholderia cepacia* complex.
2. The method of claim 1, wherein the nucleotide sequence information for bacteria of the *Burkholderia cepacia* complex in the sample and in the standard library are obtained by evaluation of restriction fragment length polymorphism.
3. The method of claim 2, wherein the restriction fragment polymorphism is carried out using the restriction enzyme *HaeIII* or *AhaI*.
4. The method of any of claims 1 to 3, wherein the *recA* gene in the bacteria of the *Burkholderia cepacia* complex in the sample is amplified relative to other nucleic acid polymers in the sample prior to obtaining the nucleotide sequence information.
5. The method of claim 4, wherein the *recA* gene is amplified using PCR amplification.
6. The method of claim 5, wherein the PCR amplification is carried out using the following primers:
Forward Primer
TGACCGCCGAGAAGAGCAA
Reverse Primer

SEQ ID No. 3

EPO - Munich
33

02 Sep. 2000

- 22 -

CTCTTCTTCGTCCATCGCCTC.

SEQ ID No. 4

7. The method of claim 5, wherein the PCR amplification is carried out using the following primers:

5 Forward Primer

TGCGGATGGGCGACGGCG

SEQ ID No. 20

Reverse Primer

CAGTTCTGTCGCTTGATCG.

SEQ ID No. 21

8. A composition comprising a pair of polynucleotide primers for production of a diagnostic amplicon from the *recA* gene of bacteria that is a member of the *Burkholderia cepacia* complex, said pair of primers hybridizing with each of the polynucleotides whose sequences are given by Seq. ID. Nos. 1, 2 and 5-19 to produce as an amplification product a diagnostic amplicon which can provide diagnostic information concerning the member of *Burkholderia cepacia* complex.

9. The composition of claim 8, wherein the polynucleotide primers have the sequences:

Forward Primer

TGACCGCCGAGAAGAGCAA

SEQ ID No. 3

Reverse Primer

CTCTTCTTCGTCCATCGCCTC.

SEQ ID No. 4

10. The composition of claim 8, wherein the polynucleotide primers have the sequence:

Forward Primer

TGCGGATGGGCGACGGCG

SEQ ID No. 20

Reverse Primer

CAGTTCTGTCGCTTGATCG.

SEQ ID No. 21

EPO - Munich
33
02. Sep. 2000

- 23 -

11. A kit for speciation of bacteria of the *Burkholderia cepacia* complex, comprising, in packaged combination, a pair of polynucleotide primers in accordance with any of claims 8 - 10, and a discriminatory restriction endonuclease.

12. The kit of claim 11, wherein the restriction endonuclease is *HaeIII* or *AluI*.

13. A composition comprising a genomovar-specific primer pair effective under stringent PCR conditions to produce amplification products by amplification of at least a portion of the *recA* gene of bacteria belonging to one genomovar of the *B. cepacia* complex, but not to produce amplification products from bacteria belonging to other genomovars.

14. The composition according to claim 13, wherein the genomovar-specific primer pairs are selected from among the following primer pairs given by Seq ID Nos.: 23 and 24, 25 and 26, 27 and 28, 29 and 30, 31 and 32, or 33 and 34.

15. A kit for speciation of bacteria of the *Burkholderia cepacia* complex, comprising, in packaged combination, a pair of genomovar-specific polynucleotide primers in accordance with claims 13 or 14 and a discriminatory restriction endonuclease.

16. The kit of claim 15, wherein the restriction endonuclease is *HaeIII* or *AluI*.

17. A vaccine composition for treatment and prevention of infection with bacteria of the *Burkholderia cepacia* complex, wherein the bacteria is a member of genomovar III and has a nucleotide sequence for the *recA* gene which produces a G-type RFLP pattern when analyzed with the restriction enzyme *HaeIII*, and wherein the vaccine composition comprises flagellin or a flagellin-derived antigen or a polynucleotide encoding flagellin or a flagellin-derived antigen, said flagellin or flagellin-derived antigen being obtained from the bacteria that is a member of genomovar III and that has a nucleotide sequence for the *recA* gene which produces a G-type RFLP pattern when analyzed with the restriction enzyme *HaeIII*.




FIG. 1

2/3




FIG. 2

Design PCR primers to conserved regions of the recA gene

Hae III digest





FIG. 3A



A B C D E F G H I J
Detect genomovar specific
restriction fragment length
polymorphisms

FIG. 3B

**COMBINED DECLARATION
AND POWER OF ATTORNEY**

As a below named inventor, I hereby declare that:

My citizenship, residence and post office address are as listed below next to my name.

I believe I am the original, first and ☒ sole/☐ joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled: **Method for the Identification and Speciation of Bacteria of the Burkholderia Cepacia Complex**

the specification of which

(a) ☒ is attached hereto.

(b) ☐ was filed on _____ as Application Serial No. _____ and was amended on _____.

(c) ☐ was described and claimed in International Application No. PCT/CA99/00813, filed on March 9, 1999 and amended on _____.

Acknowledgment of Duty of Disclosure

I hereby state that I have reviewed and understood the content of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the patentability of the subject matter claimed in this application in accordance with Title 37, Code of Federal Regulations § 1.56(a).

35 U.S.C. § 120

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) or 365(c) of any PCT International application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose material information as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

(Application Serial No.)	(Filing Date)	(Status)(patented,pending,abandoned)	(Patent No. if applicable)
--------------------------	---------------	--------------------------------------	----------------------------

(Application Serial No.)	(Filing Date)	(Status)(patented,pending,abandoned)	(Patent No. if applicable)
--------------------------	---------------	--------------------------------------	----------------------------

Power of Attorney

I hereby appoint Cari Oppedahl, PTO Reg. NO. 32,746, Marina T. Larson, PTO Reg. No. 32,038, and D'Arcy Straub, PTO Reg. No. 47,110, of the firm of OPPEDAHL & LARSON LLP, having office at P.O. Box 5068, Dillon, CO 80435-5068 as attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

SEND CORRESPONDENCE TO:



021121

PATENT TRADEMARK OFFICE

DIRECT TELEPHONE CALLS TO:
OPPEDAHL & LARSON LLP
(970)468-6800

Claim for Priority

I hereby claim foreign priority benefits under 35 U.S.C. § 119 (a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International application which designated at least one country other than the United States of America, listed below and have also identified below any foreign applications for patent or inventor's certificate, or of any PCT International application having a filing date before that of the application on which priority is claimed.


EARLIEST FOREIGN APPLICATION(S), FILED WITHIN TWELVE MONTHS (6 MONTHS FOR DESIGN) PRIOR TO SAID APPLICATION					
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)	PRIORITY CLAIMED	CERTIFIED COPY ATTACHED
				YES <input type="checkbox"/> NO <input type="checkbox"/>	YES <input type="checkbox"/> NO <input type="checkbox"/>
FOREIGN APPLICATION(S), IF ANY, FILED MORE THAN 12 MONTHS (6 MONTHS FOR DESIGN) PRIOR TO SAID APPLICATION					
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)		

Provisional Application

I hereby claim the benefit under 35 U.S.C § 119(e) of any United States provisional application(s) listed below.

<u>80/089,115</u>	<u>September 3, 1998</u>
<u>80/089,116</u>	<u>September 3, 1998</u>
(application number)	(filing date)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE OR FIRST INVENTOR <u>1 - 00</u>	LAST NAME <u>MAHENTHIRALINGAM</u>	FIRST NAME <u>ESHWAR</u>	MIDDLE NAME
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE <u>Cardiff GBN</u>	STATE OR COUNTRY OF RESIDENCE <u>Great Britain</u>	COUNTRY OF CITIZENSHIP <u>CA</u>
POST OFFICE ADDRESS <u>Cardiff School of Biosciences, Cardiff University</u> <u>P.O. Box 818,</u>		CITY <u>Cardiff, CF10 3TL, Wales</u>	STATE/COUNTRY ZIP CODE <u>Great Britain</u>
DATE <u>16 FEBRUARY 2001</u>		SIGNATURE 	

- ☐ Signature for additional joint inventor attached. Number of Pages ____.
- ☐ Signature by Administrator(trix) or legal representative for deceased or incapacitated inventor. Number of Pages ____.
- ☐ Signature for inventor who refuses to sign or cannot be reached by person authorized under 37 CFR § 1.47. Number of Pages ____.

09763298-022001

UBC.P-017
PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants: Mahenthiralingam, E.
Serial No.: 09/763,298
Filed : September 23, 1999 (International Application)
For : Method for the Identification and Speciation of Bacteria of the
Burkholderia cepacia Complex

SUBMISSION OF SEQUENCE LISTING

Sir:

Responsive to the Notice of Missing Requirements mailed March 27, 2001 for the above-captioned application, Applicants enclose a sequence diskette. The undersigned certifies that the contents of this diskette are the same as the printed sequence listing which appears in the specification as filed.

Respectfully submitted,



Marina T. Larson

Patent Office Reg. No. 32,038
Attorney for Applicants
(970) 668-6600 x152

I hereby certify that this paper is being sent by first class mail, postage pre-paid to Asst. Commissioner for Patents, Washington, D.C. 20231 on April 4, 2001.



Marina T. Larson, PTO # 32,038

April 4, 2001

Date of Signature

09/763,298-022001

WO 00/14274

SEQUENCE LISTING

<110> University of British Columbia
Mahenthiralingam, Eshwar

<120> Method for the Identification and Speciation of
Bacteria of the Burkholderia Cepacia Complex

<130> 80472-5

<140>

<141>

<150> 60/099,115

<151> 1998-09-03

<150> 60/099,116

<151> 1998-09-03

<160> 40

<170> PatentIn Ver. 2.0

<210> 1

<211> 1044

<212> DNA

<213> Burkholderia .cepacia

<220>

<223> recA gene

<400> 1

```

atgaccgcgcg agaagagcaa ggcgctggct gccgcactcg cgcagatcga gaagcagttc 60
ggcaaaggggt cgatcatgcg catgggcgac ggcgaggcgg ccgaggacat ccaggtcgtg 120
tccacggggt cgctcggggt cgacatcgca ctgggcgtcg gcggcctgcc gcgcggccgc 180
gtggtcgaga tctacgggcc ggaatcgtcc ggcaagacca cgctcacgct gcaggtcatt 240
gccgagctgc agaagctggg cggcacccgc gcgttcatcg acgccgagca cgcgctcgac 300
gtccagtagc cggcgaagct cggcgtcaac gtgccggatc tgctgatctc gcagccggac 360
accggcgagc aggcgctcga aatcacccgc gcgctggtgc gctcgggctc gatcgacatg 420
atcgatcatc actcgggtcg ggcgctcgtg ccgaaggcgg aaatcgaagg cgagatgggc 480
gattcgctgc cgggcctgca ggcgcgcctg atgtcgcagg cgctgcgcaa gctgacgggc 540
acgatcaagc gcacgaactg cctcgtgatc ttcatcaacc agatccgcat gaagatcggc 600
gtgatgttcg gcaaccgcga aaccacgacg ggcggcaacg cgctgaagtt ctattcgtcg 660
gtgctgtctc acatccgcgg gatcgggtcg atcaagaaga acgacgaggt gatcggcaac 720
gaaacgcgcg tgaaggtcgt caagaacaag gtgtcgccgc cgttcgcgca agcgatcttc 780
gacatcctgt acggcgaggg catttcgct cagggcgaga tcattgatct cggcgtgcag 840
gcgaagatcg tgcacaaggc gggcgccctg tacagctaca acggcgagaa gatcggtcag 900
ggcaaggaca acgcgcgtga attcctgcgc gagaatccgg aaatcgcgcg cgagatcgag 960
aaccgcattc gtgaatcgct cggcgttgcc gccatgccgg atggcgcggg ccacgacgaa 1020
gccgaggcga tggacgaaga agag                                     1044

```

<210> 2

<211> 1044

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 2

```

atgaccgccg agaagagcaa ggcgctggcg gccgcactcg cgcagatcga gaagcagttc 60
ggcaaagggt cgatcatgcg gatgggcgac ggcgacgtga aggaagacat ccaggctcgtc 120
tccacgggct cgctgggcct cgatatcgcg ctccggcgctg gcggcctgcc gcgcggccgg 180
gtggtcgaaa tctacggtcc ggaatcgctc ggcaagacca cgctcacgct gcagggtgatc 240
gccgagctgc agaagctggg cggcacggca gccttcacatc acgccgagca cgcgctcgac 300
gtccagtagc cggcgaagct cggcgtgaac gtgccggagc tgctgatctc gcagccggac 360
accggcgaac aggcgctcga aatcaaccgac gcgctcgtgc gctcgggctc gatcgacatg 420
atcgatcatc actcggtcgc ggcaactggg ccgaaggccg atatcgaagg cgagatgggc 480
gattcgctgc cgggcctgca tgcgcgtctg atgtcgacag cgctgcgcaa gctgacgggc 540
acgatcaagc gcacgaactg cctcgtgatc ttcacaaacc agatccgcat gaagatcggc 600
gtgatgttcg gcaaccggga aaccacgacg ggccgcaacg cgctgaagtt ctattcgtcg 660
gtgcgtctcg acatccgccc gatcggctcg atcaagaaga acgacgaggt gatcggcaac 720
gaaacgcgcg tgaaggctgt caagaacaag gtgtcgccgc cgttcgcgca agcgatcttc 780
gacatcctgt acggcgaggg catttcgcgt caggggcgaga tcacgatctc cggcgtgcag 840
gcgaagatcg tcgacaaggc gggcgccctg tacagctaca acggcgagaa gatcggccag 900
ggcaaggaca acgcgcgtga attcctgcgc gagaatccgg aaatcgcgcg tgagatcgag 960
aaccgcattc gtgaatcgct cggcgttgcc gccatgccgg atggcgcggg ccacgacgaa 1020
ccgaggcgat gggacgaaga agag                                     1044

```

<210> 3

<211> 19

<212> DNA

<213> Burkholderia cepacia

<220>

<223> B. cepacia recA forward primer (BCR1)

<400> 3

tgaccgccga gaagagcaa

19

<210> 4

<211> 21

<212> DNA

<213> Burkholderia cepacia

<220>

<223> B. cepacia recA reverse primer (BCR2)

<400> 4

ctcttcttcg tccatcgct c

21

<210> 5

<211> 1044

<212> DNA

<213> Burkholderia multivorans

<220>

<223> recA gene

<400> 5

```

atgaccgccg agaagagcaa ggcgctggct gccgcactcg cgcagatcga gaagcagttc 60
ggcaaagggg cgatcatgcg catgggcgac ggcgaggcgg ccgaggacat ccaggtcgtg 120
tccacggggt cgctcgggct cgacatcgca ctgggctgct ggcgcctgcc gcgcggccgc 180
gtggtcgaga tctacggggt ggaatcgctc ggcaagacca cgctcacgct gcaggtcatc 240
gccgagctgc agaagctggg cggcactgct gcgttcacgc acgccgagca cgcgctcgac 300
gtccagtagc cggcgaagct cggcgtaaac gtgccggatc tgctgatctc gcagccggac 360
accggcgagc aggcgctcga aatcacgcac gcgtcgtgct gctcgggctc gatcgacatg 420
atcgctcatc actcggctcg ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggcctgca ggcgcgcctg atgtcgagc cgctgcgcaa gctgacgggc 540
acgatcaagc gcacgaactg cctcgtgatc ttcataaacc agatccgcat gaagatcggc 600
gtgatgttcg gcaaccggga aaccacgacg gccggcaacg cgctgaagtt ctattcgtcg 660
gtgcgtctcg acatccggcg gatcggctcg atcaagaaga acgacgaggt gatcggcaac 720
gaaacgcgcg tgaaggtcgt caagaacaag gtgtcgccgc cgttccgcga agcgatcttc 780
gacatcctgt acggcgaggg catttcgcgt cagggcgaga tcatcgatct cggcgtgcag 840
gcgaagatcg tcgacaaggc ggcgcgcctg tacagctaca acggcgagaa gatcggtcag 900
ggcaaggaca acgcgcgtga attcctgcgc gagaatccgg aaatcgcgcg cgagatcgag 960
aaccgcattc gtgaatcgct cggcgttgcc gccatgccgg atggcgcggg ccacgacgaa 1020
gccgaggcga tggacgaaga agag                                     1044

```

<210> 6

<211> 1044

<212> DNA

<213> Burkholderia multivorans

<220>

<223> recA gene

<400> 6

```

atgaccgccg agaagagcaa ggcgctggct gccgcactcg cgcagatcga gaagcagttc 60
ggcaaagggg cgatcatgcg catgggcgac ggcgaggcgg ccgaggacat ccaggtcgtg 120
tccacggggt cgctcgggct cgacatcgca ctgggctgct ggcgcctgcc gcgcggccgc 180
gtggtcgaga tctacggggt ggaatcgctc ggcaagacca cgctcacgct gcaggtcatc 240
gccgagctgc agaagctggg cggcaccgca gcgttcacgc acgccgagca cgcgctcgac 300
gtccagtagc cggcgaagct cggcgtaaac gtgccggatc tgctgatctc gcagccggac 360
acgggcgagc aggcgctcga aatcacgcac gcgtcgtgct gctcgggctc gatcgacatg 420
atcgctcatc actcggctcg ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggcctgca ggcgcgcctg atgtcgagc cgctgcgcaa gctgacgggc 540
acgatcaagc gcacgaactg cctcgtgatc ttcataaacc agatccgcat gaagatcggc 600
gtgatgttcg gcaaccggga aaccacgacg gccggcaacg cgctgaagtt ctattcgtcg 660
gtgcgtctcg acattcggcg gatcggctcg atcaagaaga acgacgaggt gatcggcaac 720
gaaacgcgcg tgaaggtcgt caagaacaag gtgtcgccgc cgttccgcga agcgatcttc 780
gacatcctgt acggcgaggg catttcgcgt cagggcgaga tcatcgatct cggcgtgcag 840
gcgaagatcg tcgacaaggc ggcgcgcctg tacagctaca acggcgagaa gatcggccag 900
ggcaaggaca acgcgcgtga attcctgcgc gagaatccgg aaatcgcgcg cgagatcgag 960
aaccgcattc gtgaatcgct cggcgttgcc gccatgccgg atggcgcggg ccacgacgaa 1020
gccgaggcga tggacgaaga agag                                     1044

```

<210> 7

<211> 1044

<212> DNA

<213> Burkholderia multivorans

<220>

<223> recA gene

<400> 7

```

atgaccgccg agaagagcaa ggcgctggct gccgcactcg cgcagatcga gaagcagttc 60
ggcaaagggt cgatcatgcy catgggcyac ggcgagggcg ccaggacat ccaggctcgtg 120
tccacggggt cgctcgggct cgacatcgca ctgggcgtcg gcggcctgcc gcgcggcccg 180
gtggtcgaga tctacgggct ggaatcgctc ggcaagacca cgtcacgct gcagggtcatc 240
gccgagctgc agaagctggg cggtaaccga gcgttcacat acgcgcagca cgcgctcgac 300
gtccagtagc cggcgaaagt cggcgtaaac gtgcgggagc tgcgtgatct tcagccggac 360
accggcgagc aggcgctcga aatcacgcac gcgctgggtg gctcgggctc gatcgacatg 420
atcgatcatc actcggctcg ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggcctgca ggcgcgcctg atgtgcgagg cgtgcgcgaa gctgacgggc 540
acgatcaagc gcacgaactg cctcgtgatc ttcataaacc agatccgcat gaagatcggc 600
gtgatgttcg gcaaccgcga aaccacgacg ggcggcaacg cgtgaagtt ctattcgtcg 660
gtgcgtctcg acatccgcgc gatcggctcg atcaagaaga acgacgaagt gatcggcaac 720
gaaaccgcgc tgaaggctcg caagaacaag gtgtcgccgc cgttcgcgca agcagatcttc 780
gacatcctgt acggcgaggc catttcgcgt cagggcgaga tcatcgatct cggcggtgcag 840
gcgaagatcg tcgacaaggc ggcgcgcctg tacagctaca acggcgagaa gatcgggtcag 900
ggcaaggaca acgcgcgtga attcctgcgc gagaatcccg aaatcgcgcg cgagatcgag 960
aaccgcattc gtgaatcgct cggcggttgc gccatgcccg atggcgcggg ccacgacgaa 1020
gccgaggcga tggacgaaga agag                                     1044

```

<210> 8

<211> 1041

<212> DNA

<213> Burkholderia vietnamiensis

<220>

<223> recA gene

<400> 8

```

atgaccgccg agaagagcaa ggcgctggcg gccgcactcg cgcagatcga gaagcagttc 60
ggcaaagggt cgatcatgcy gatgggcyac ggcgacgtga aggaagacat ccaggctcgtc 120
tccacggggt cgctcgggct cgatatacgc ctccggctcg gcggcctgcc gcgcggcccg 180
gtggtcgaaa tctacgggct ggaatcgctc ggcaagacca cgtcacgct gcagggtgatc 240
gccgagctgc agaagctggg cggcacggca gccttcacat acgcgcagca cgcgctcgac 300
gtccagtagc cggcgaaagt cggcgtaaac gtgcgggagc tgcgtgatct gcagccggac 360
accggcgaaac aggcgctcga aatcacgcac gcgctcgtgc gctcgggctc gatcgacatg 420
atcgatcatc actcggctcg ggcactgggt ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggtctgca ggcgcgtctg atgtcgcagg cgtgcgcgaa gctgacgggc 540
acgatcaagc gcacgaactg cctcgtgatc ttcataaacc agatccggat gaagatcggc 600
gtgatgttcg gcaaccgcga aaccacgacg ggcggcaatg cactgaagtt ctattcgtcg 660
gtgcgtctcg acatccgcgc gatcggctcg atcaagaaga acgacgaggt gatcggcaac 720
gaaaccgcgc tgaaggctcg caagaacaag gtatcgccgc cgttcgcgca agcagatcttc 780
gacatcctct acggcgaggc catttcgcgt cagggcgaga tcatcgatct cggcggtgcag 840
gcgaaaatcg tcgacaaggc cggcgccctg tacagctaca acggcgagaa gatcgggtcag 900
ggcaaggaca acgcgcgtga attcctgcgc gagaatcccg aaatcgcgcg cgagatcgag 960
aaccgcattc gcgagtcgct cgggtgctgc gcgatgcccg atggcggtgt cgacgaagcc 1020
gaggcgatgg acgaagaaga g                                     1041

```

<210> 9
 <211> 1041
 <212> DNA
 <213> Burkholderia cepacia

<220>
 <223> recA gene

<400> 9
 atgaccgccc agaagagcaa ggcgcttgcc gccgcactcg cgcagatcga gaagcagttc 60
 ggcaaagggc cgatcatgcg catggggcgac ggcgaggcgg cggaggatat ccaggctcgtc 120
 tccacgggct cgctgggtct cgacatcgca ctggggcgctg gcggcctgcc gcgcggccgg 180
 gtggctgaaa tctacggtec ggaatcgtec ggtaaaacca cgctcacgct gcaggctcatc 240
 gccgaactgc agaagatcgg cggcacggct gcgttcacgc acgccgaaca cgcgctcgac 300
 gttcagtagc cgtcgaagct cggcgtgaac gtgccggaac tgctgatctc gcagccggac 360
 accggcgagc aggcgcttga aatcacgcac gcgctggtgc gctcgggctc gatcgacatg 420
 atcgatcatc actcggctcg ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggc 480
 gattcgctgc cgggcctgca ggcgcgcctg atgtcgagg cgctgcgcaa gctgaccggc 540
 acgatcaagc gcacgaactg cctcgtgatc ttcatacaac agatccggat gaagatcggc 600
 gtgatgttcg gcaacccgga aaccacgacg ggccgcaacg cgctgaagtt ctatgcgtcg 660
 gtgcgccttg atatccgccg gatcggctcg atcaagaaga acgacgaggt gatcggcaac 720
 gaaacccgcg tgaaggtcgt caagaacaag gtatcgccgc cgttcgcgca agcgatcttc 780
 gacatcctgt atggcgaggg catttcgcgc cagggcgaga tcatcgatct cggcgtgcag 840
 gcgaagatcg tcgacaaggc gggcgccctg tacagctaca acggcgagaa gatcggccag 900
 ggcaaggaca acgcgcgtga attcctgcgc gagaatccgg aaatcgcacg cgagatcgag 960
 aaccgcattc gcgaatcgct cgggtgctgt gccatgcccc atggcgtagt caacgaagcc 1020
 gaggcgatgg acgaagaaga g 1041

<210> 10
 <211> 1041
 <212> DNA
 <213> Burkholderia cepacia

<220>
 <223> recA gene

<400> 10
 atgaccgccc agaagagcaa ggcgcttgcc gccgcactcg cgcagatcga gaagcagttc 60
 ggcaaagggc cgatcatgcg catggggcgac ggcgaggcgg cggaggatat ccaggctcgtc 120
 tccacgggct cgctgggtct cgacatcgca ctggggcgctg gcggcctgcc gcgcggccgg 180
 gtggctgaaa tctacggtec ggaatcgtec ggtaaaacca cgctcacgct gcaggctcatc 240
 gccgaactgc agaagatcgg cggcacggct gcgttcacgc acgccgaaca cgcgctcgac 300
 gttcagtagc cgtcgaagct cggcgtgaac gtgccggaac tgctgatctc gcagccggac 360
 accggcgagc aggcgcttga aatcacgcac gcgctggtgc gctcgggctc gatcgacatg 420
 atcgatcatc actcggctcg ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggc 480
 gattcgctgc cgggcctgca ggcgcgcctg atgtcgagg cgctgcgcaa gctgaccggc 540
 acgatcaagc gcacgaactg cctcgtgatc ttcatacaac agatccggat gaagatcggc 600
 gtgatgttcg gcaacccgga aaccacgacg ggccgcaacg cgctgaagtt ctatgcgtcg 660
 gtgcgccttg atatccgccg gatcggctcg atcaagaara acgacgaggt gatcggcaac 720
 gaaacccgcg tgaaggtcgt caagaacaag gtatcgccgc cgttcgcgca agcgatcttc 780
 gacatcctgt atggcgaggg catttcgcgc cagggcgaga tcatcgatct cggcgtgcag 840
 gcgaagatcg tcgacaaggc gggcgccctg tacagctaca acggcgagaa gatcggccag 900

```

ggcaaggaca acgcgcgtga attcctgcgc gagaatccgg aaatcgcacg cgagatcgag 960
aaccgcattc gcgaatcgct cgggtgctgt gccatgcccc atggcgtagt caacgaagcc 1020
gaggcgatgg acgaagaaga g                                     1041

```

<210> 11

<211> 1041

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 11

```

atgaccgccc agaagagcaa ggcgctggcg gccgcactcg cacagatcga gaagcagttc 60
ggcaaggggt cgatcatgcg catgggcgac ggcgaggcgg ccgagaatat ccaggtcgtc 120
tccacggggg cgctgggtct cgacatcgcg ctgggcgtcg gcggtttgcc gcgcggccgg 180
gtggtcgaga tctacggtcc ggaatcgctc ggtaaaacca ccctcacgct gcaagtcatc 240
gctgaactgc agaagatcgg cggcacggca gccttcacg acgccgagca tgcactcgac 300
gtccagtagc cgtcgaagct cggcggtgaac gtgccggaac tgctgatctc gcagccggac 360
accggcgagc aggcgctcga aatcacccgat gcgctggtgc gctcgggctc gatcgacatg 420
atcgatcatc actcggtcgc ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggt 480
gattcgctgc cgggcctgca ggctcgctcg atgtcgcagg cgctgcgcaa gctgaccggc 540
acgatcaagc gcacgaactg cctcgtgatc ttcatcaacc agatccggat gaagatcggc 600
gtgatgttcg gcaaccggga aaccaccacg ggcggaacg cgctgaagtt ctattcgctc 660
gtgctgtctc acattcgccg gatcggctcg atcaagaaga acgacgaggt gatcggcaac 720
gaaaccccgc tgaaggctgt caagaacaag gtgtcgccgc cgttcgcgca agcgatcttc 780
gacatcctgt acggcggaagg catttcgctg cagggcgaga tcatcgatct cggcggtgcg 840
gcgaagatcg tcgacaaggc cggcgctcgt tacagctaca acggcgagaa gatcggccag 900
ggcaaggaca atgcgcgtga attcctgcgc gagaatccgg aaatcgcacg cgagatcgag 960
aaccgcattc gcgaatcgct cgggtgctgt agcatgcccc atggcgtagc caacgaagcc 1020
gaggcgatgg acgaagaaga g                                     1041

```

<210> 12

<211> 1041

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 12

```

atgaccgccc agaagagcaa ggcgctggcg gccgcactcg cacagatcga gaagcagttc 60
ggcaaggggt cgatcatgcg catgggcgac ggcgaggcgg ccgagaatat ccaggtcgtc 120
tccacggggg cgctgggtct cgacatcgcg ctgggcgtcg gcggcctgcc gcgcggccgg 180
gtggtcgaga tctacggtcc ggaatcgctc ggtaaaacca ccctcacgct gcaagtcatc 240
gctgaactgc agaagatcgg cggcacggca gccttcacg acgccgagca cgctcgcgac 300
gtccagtagc cgtcgaagct cggcggtgaac gtgccggaac tgctgatctc gcagccggac 360
accggcgagc aggcgctcga aatcacccgat gcgctggtgc gctcgggctc gatcgacatg 420
atcgatcatc actcggtcgc ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggtctgca ggctcgctcg atgtcgcagg cgctgcgcaa gctgaccggc 540
acgatcaagc gcacgaactg cctcgtgatc ttcatcaacc agatccggat gaagatcggc 600
gtgatgttcg gcaaccggga aaccaccacg ggcggaacg cgctgaagtt ctattcgctc 660
gtgctgtctc acattcgccg gatcggctcg atcaagaaga acgatgaggt gatcggcaac 720

```

```

gaaacccgcg tgaaggctcg caagaacaag gtgtcgccgc cgttccgcga agcgatcttc 780
gacatcctgt acggcgaagg catttcgcgt cagggcgaga tcatcgatct cggcggtgcag 840
gcgaagatcg tcgacaaggc gggcgccctg tacagctaca acggcgagaa gatcggccag 900
ggcaaggaca atgcgcgtga attcctgcgt gagaatccgg aaatcgcacg cgagatcgaa 960
aaccgcatcc gcgaatcgct cgggtgtcgt agcatgcccg atggcgtagc caacgaagcc 1020
gagggcgatgg acgaagaaga g                                     1041

```

<210> 13

<211> 1041

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 13

```

atgaccgccc agaagagcaa ggcgctggcg gccgcactcg cgcagatcga gaagcagttc 60
ggcaagggct cgatcatgct catgggagac ggcgaggcga ccgagaatat ccaggctcgtc 120
tccacggggt cgctgggtct cgacatcgct ctgggcgttg gcggcctgcc gcgcggcccg 180
gtggtcgaga tctacggtcc ggaatcgtcc ggtaagacca cctcacgct gcaagtcattc 240
gctgaactgc agaagatcgg cggcacggca gccttcacg acgcagagca cgcgctcgac 300
gtccagtagc cgtcgaagct cggcgtgaac gtgccgggaa tgotgatctc gcagccggac 360
accggtagc aggcgctcga aatcacgcat gcgctgggtg gctcgggctc gatcgacatg 420
atcgatcatg actcggctcg ggcgctcgt ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggtctgca ggcccgcctg atgtcgcagg cgtcgcgcaa gctgaccggc 540
acgatcaagc gcacgaactg cctgggtgatc ttcacaaacc agatccggat gaagatcggc 600
gtgatgttcg gcaaccggga aaccaccacg ggcggaacg cgtcgaagtt ctattcgtcg 660
gtgcgtctcg acattcgccg gatcggctcg atcaagaaga acgacgaggt gatcggaac 720
gaaacccgcg tgaaggctcg caagaacaag gtgtcgccgc cgttccgcga agcgatcttc 780
gacatcctgt acggcgaagg tatttcgcgt cagggcgaga tcatcgatct cggcggtgcag 840
gcgaagatcg tcgacaaggc cggcgccctg tacagctaca acggcgagaa gatcggccag 900
ggcaaggaca atgcgcgtga attcctgcgc gagaatccgg aaatcgcacg cgagatcgaa 960
aaccgcatcc gcgaatcgct cgggtgtcgt agcatgcccg atggcgtagc caacgaagcc 1020
gagggcgatgg acgaagaaga g                                     1041

```

<210> 14

<211> 1041

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 14

```

atgaccgccc agaagagcaa ggcgctggcc gccgcgcttg ccagatcga gaagcagttc 60
ggcaagggct cgatcatgct catgggagac ggcgaggcgg ccgaagatat ccaggctcgtc 120
tccacggggt cgctgggctt cgatcgcgct cttggcgctg gcggcctgcc gcgcggcccg 180
gtggtcgaga tctacggccc ggaatcgtcc ggtaaaacca cgtcacgct gcaggtcatt 240
gccgagctgc agaagctggg cggcacggca gcgttcacg acgcagagca cgcgctcgac 300
gtccagtagc cgtcgaagct cggcgtgaat gtgccggagc tgotgatttc gcaaccggac 360
accggcgagc aggcgctgga aatcacgcat gcgctgggtg gctcgggctc gatcgacatg 420
atcgatcatg actcggctcg ggcgctcgt ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggcctgca ggcccgcctg atgtcgcagg cgtcgcgcaa gctgaccggc 540

```



```

acgatcaagc gcacgaactg cctggtgata ttcatacaacc agattcgtat gaagatcggc 600
gtgatgttcg gcaacccgga aaccacgacg ggcggtaacg cgctgaagtt ctatgcgtcg 660
gtgctgtctc atatccgccg gatcggctcg atcaagaaga acgacgaggt gatcggcaac 720
gaaacccgtg tgaaggctgt caagaacaag gtgtcgccgc cgttccgcga agcgatcttc 780
gacatcctgt atggcgaggg catttcgcgt caggcgagaga tcatcgatct cggcgtgcag 840
gcgaagatcg tcgacaaggc aggcgcctgg tacagctaca acggcgagaa gattggccag 900
ggcaaggaca acgcgcgtga attcctgcgc gagaatccgg aaatcgcgcg cgagatcgag 960
aaccgcatcc gcgaatcgct cggcgtcgtc gcaatgcccg atggtgcagg caacgaagcc 1020
gaggcgatgg acgaagaaga g                                     1041

```

<210> 15

<211> 1041

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 15

```

atgaccgccg agaagagcaa ggcgctggcc gcgcgcgttg ccagatcga gaagcagttc 60
ggcaagggct cgatcatgcg catggggcag ggcgagggcg cggagatat ccaggtcgtc 120
tccacggggt cgctgggcct cgatatcgcg cttggcgctg gcggcctgcc gcgcggcccg 180
gtggtcgaga tctacggccc ggaatcgctc ggtaaaacca cgctcacgct gcaggtcatc 240
gccgagctgc agaagctggg cggcaccgca gcgttcacg acgcgagca cgcgctcgac 300
gtccagtagc cgtcgaaagt cggcgtgaat gtgccggagc tgctgatttc gcagccggac 360
accggcgagc aggcgctgga aatcaccgat gcgctggtgc gctcgggctc gatcgacatg 420
atcgtcatcg actcggctgc ggcgctcgct ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggcctgca ggcccgcttg atgtcgcagg cgctgcgcaa gctgaccggc 540
acgatcaagc gcacgaactg cctggtgata ttcatacaacc agattcgtat gaagatcggc 600
gtgatgttcg gcaacccgga aaccacgacg ggcggtaacg cgctgaagtt ctatgcgtcg 660
gtgctgtctc atatccgccg gatcggctcg atcaagaaga acgacgaggt gatcggcaac 720
gaaacccgtg tgaaggctgt caagaacaag gtgtcgccgc cgttccgcga agcgatcttc 780
gacatcctgt atggcgaggg catttcgcgt caggcgagaga tcatcgatct cggcgtgcag 840
gcgaagatcg tcgacaaggc aggcgcctgg tacagctaca acggcgagaa gattggccag 900
ggcaaggaca acgcgcgtga attcctgcgc gagaatccgg aaatcgcgcg cgagatcgag 960
aaccgcatcc gcgaatcgct cggcgtcgtc gcaatgcccg atggtgcagg caacgaagcc 1020
gaggcgatgg acgaagaaga g                                     1041

```

<210> 16

<211> 1041

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 16

```

atgaccgccg agaagagcaa ggcgctggcg gcgcgcgtcg ccagatcga aaagcagttc 60
ggcaagggct cgatcatgcg gatggggcag ggcgaagcgg ccgaggatat ccaggtcgtc 120
tccacggggt cgctgggtct cgacatcgcg ctgggcgtcg gcggcttgcc gcgcggcccg 180
gtggtcgaga tctacggctc ggaatcgctc ggtaaagacca cgctcacgct gcaggtcatc 240
gccgagctgc agaagctggg cggcaccgcg gcgttcacg acgcgagca cgcgctcgac 300
gttcaatatg ccgcgaagct cggcgtgaac gtgccggagc tgctgatctc gcagccggac 360

```

```

accggcgagc aggccctcga aatcacccgat gcgctgggtgc gctcggggctc gatcgacatg 420
atcgctcatcg actcgggtcgc ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggtctgca ggcccgcctg atgtcgcagg cgctgcgcaa gctgaccggt 540
acgatcaagc gcacgaactg cctcgtgacg ttcacaaacc agatccggat gaagatcggc 600
gtgatgttcg gcaacccgga aaccacgacg ggcggtaacg cactgaagtt ctactcgtcg 660
gtgctctcgc atatccgccg gattggctcg atcaagaaga gcgacgaggt gatcggcaac 720
gaaacccgcg tgaaggtcgt caagaacaag gtgtcgccgc cgttccgcga agcgatcttc 780
gacatcctgt acggcggaagg catttcgcgt cagggcgaga tcatcgatct cggcgtgcag 840
gcgaagatcg tcgacaaggc gggcgccctg tacagctaca acggcgagaa gatcggccag 900
ggcaaggaca acgcgcgcga attcctgcgc gaaaatccgg aaatcgcgcg cgagatcgag 960
aaccgcatcc gcgaatcgct cggcgtcgtc gcaatgcccg atggcgcagg caacgaagcc 1020
gaggcgatgg acgaagaaga g                                     1041

```

<210> 17

<211> 1041

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 17

```

atgaccgcgc agaagagcaa ggcgctggcg gccgcgctcg cgcagatcga aaagcagttc 60
ggcaagggct cgatcatgcg gatgggcgac ggcggaagcgg ccgaggatat ccaggtcgtc 120
tcacgggggt cgctgggtct cgacatcgcg ctgggcgtcg gcggcttgcc gcgcggcccg 180
gtggctcgaga tctacgggtc ggaatcgtcc ggtaagacca cgctcacgct gcaggtcatc 240
gccgagctgc agaagctggg cggcaccgcg gcgttcacg acgcccagca cgcgctcgac 300
gttcaatatg ccgcgaagct cggcgtgaac gtgccggagc tgetgatctc gcagccggac 360
accggcgagc aggccctcga aatcacccgat gcgctgggtgc gctcggggctc gatcgacatg 420
atcgctcatcg actcgggtcgc ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgctgc cgggtctgca ggcccgcctg atgtcgcagg cgctgcgcaa gctgaccggt 540
acgatcaagc gcacgaactg cctcgtgacg ttcacaaacc agatccggat gaagatcggc 600
gtgatgttcg gcaacccgga aaccacgacg ggcggtaacg cactgaagtt ctactcgtcg 660
gtgctctcgc atatccgccg gatcggctcg atcaagaaga acgacgaggt gatcggcaac 720
gaaacccgcg tgaaggtcgt caagaacaag gtgtcgccgc cgttccgcga agcgatcttc 780
gacatcctgt acggcggaagg catttcgcgt cagggcgaga tcatcgatct cggcgtgcag 840
gcgaagatcg tcgacaaggc gggcgccctg tacagctaca acggcgagaa gatcggccag 900
ggcaaggaca acgcgcgcga attcctgcgc gaaaatccgg aaatcgcgcg cgagatcgag 960
aaccgcatcc gcgaatcgct cggcgtcgtc gcaatgcccg atggcgcagg caacgaagcc 1020
gaggcgatgg acgaagaaga g                                     1041

```

<210> 18

<211> 1041

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 18

```

atgaccgcgc agaagagcaa agcgctggcg gccgcgctcg cgcagatcga aaagcagttc 60
ggcaagggct cgatcatgcg gatgggcgac ggcggaagcgg ccgaggatat ccaggtcgtc 120
tcacgggggt cgctgggtct cgacatcgcg ctgggcgtcg gcggcttgcc gcgcggcccg 180

```

```

gtgggtcgaga tctacgggtcc ggaatcgtcc ggtaagacca cgctcacgct gcaggtcatc 240
gccgagctgc agaagctggg cggcaccgcg gcgttcatcg acgccgagca cgcgctcgac 300
gttcaatatg ccgcgaagct cggcgtgaac gtgccggagc tgctgatctc gcagccggac 360
accggcgagc agggcctcga aatcaccgat gcgctgggtgc gctcgggctc gatcgacatg 420
atcgatcatcg actcgggtcgc ggcgctcgtg ccgaaggccg aaatcgaagg cgagatgggc 480
gattcgtctgc cgggtctgca ggcccgcctg atgtcgcagg cgctgcgcaa gctgaccggg 540
acgatcaagc gcacgaactg cctcgtgatc ttcatcaacc agatccggat gaagatcggc 600
gtgatgttcg gcaacccgga aaccacgacg ggcggttagc cactgaagtt ctactcgtcg 660
gtgctgtctc atattccgcg gatcgggtcgc atcaagaaga acgacgaggt gatcggcaac 720
gaaacccgcg tgaaggctcg caagaacaag gtgtcgcgcg cgttcgcgca agcgtcttc 780
gacatcctgt acggcgaagg catttcgcgt cagggcgaga tcatcgatct cggcgtgcag 840
gcgaagatcg tcgacaaggc gggcgcttg tacagctaca acggcgagaa gatcggccag 900
ggcaaggaca acgcgcgcga attcctgcgc gaaaatccgg aaatcgcgcg cgagatcgag 960
aaccgcatcc gcgaatcgtc cggcgtcgtc gcaatgcccg atggcgcagg caacgaagcc 1020
gaggcgatgg acgaagaaga g                                     1041

```

<210> 19

<211> 1041

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA gene

<400> 19

```

atgaccgccc agaagagcaa ggcgctggcg gccgcgctcg cgcagatcga aaagcagttc 60
ggcaagggct cgatcatgcy gatgggcgac ggcaagcgg ccgaggatat ccaggtcgtc 120
tccacggggc cgctgggtct cgacatcgcy ctgggcgtcg gcggttgcc gcgcggccgg 180
gtgggtcgaga tctacgggtcc ggaatcgtcc ggtaagacca cgctcacgct gcaggtcatc 240
gccgagctgc agaagctggg cggcaccgcy gcgttcatcg acgccgagca cgcgctcgac 300
gttcaatatg ccgcgaagct cggcgtgaac gtgccggagc tgctgatctc gcagccggac 360
accggcgaac atgccctcga aatcaccgat gcgctgggtgc gctcgggctc gatcgacatg 420
atcgatcatcg actcgggtcgc ggcgctcgtg ccgaaaggcc gaaatcgaat gcgaaatggg 480
cgattcgtct cgggtctgca agggcccctg atgtcgcagg cgctgcgcaa gctgaccggg 540
acgatcaagc gcacgaactg cctcgtgatc ttcatcaacc agatccggat gaagatcggc 600
gtgatgttcg gcaacccgga aaccacgacg ggcggtaacg cactgaagtt ctactcgtcg 660
gtgctgtctc atattccgcg gattgggtcg atcaagaaga gcgacgaggt gatcggcaac 720
gaaacccgcy tgaaggctcg caagaacaag gtgtcgcgcg cgttcgcgca agcgtcttc 780
gacatcctgt acggcgaagg catttcgcgt cagggcgaga tcatcgatct cggcgtgcag 840
gcgaagatcg tcgacaaggc gggcgcttg tacagctaca acggcgagaa gatcggccag 900
ggcaaggaca acgcgcgcga attcctgcgc gaaaatccgg aaatcgcgcg cgagatcgag 960
aaccgcatcc gcgaatcgtc cggcgtcgtc ccaatgcccg atggcgcagg caacgaagcc 1020
gaggcgatgg acgaagaaga g                                     1041

```

<210> 20

<211> 18

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA forward primer (BCRU1*)

<400> 20

tgccgatggg cgacggcg

18

<210> 21

<211> 19

<212> DNA

<213> Burkholderia cepacia

<220>

<223> recA reverse primer (BCRU2*)

<400> 21

cagttctgtc gcttgatcg

19

<210> 22

<211> 19

<212> DNA

<213> Burkholderia multivorans

<220>

<223> B. multivorans specific recA forward primer
(BCRBM1)

<400> 22

cggcgtcaac gtgccggat

19

<210> 23

<211> 19

<212> DNA

<213> Burkholderia multivorans

<220>

<223> B. multivorans specific reA reverse primer
(BCRBM2)

<400> 23

tccatcgct cggettcgt

19

<210> 24

<211> 18

<212> DNA

<213> Burkholderia vietnamiensis

<220>

<223> B. vietnamiensis specific recA forward primer
(BCRBV1)

<400> 24

gggcgacggc gacgtgaa

18

<210> 25

<211> 18

<212> DNA

<213> Burkholderia vietnamiensis

<220>
 <223> B. vietnamiensis specific recA reverse primer
 (BCRBV2)

<400> 25
 tcggccttcg gcaccagt 18

<210> 26
 <211> 18
 <212> DNA
 <213> Burkholderia cepacia

<220>
 <223> B. cepacia Genomovar IV specific recA forward
 primer (BCRG41)

<400> 26
 accggcgagc aggcgctt 18

<210> 27
 <211> 18
 <212> DNA
 <213> Burkholderia cepacia

<220>
 <223> B. cepacia Genomovar IV specific recA reverse
 primer (BCRG42)

<400> 27
 acgccatcgg gcatggca 18

<210> 28
 <211> 18
 <212> DNA
 <213> Burkholderia cepacia

<220>
 <223> B. cepacia Genomovar III, RG-B recA specific
 forward primer

<400> 28
 gcaagtcac gctgagaa 18

<210> 29
 <211> 18
 <212> DNA
 <213> Burkholderia cepacia

<220>
 <223> B. cepacia Genomovar III, RG-B recA specific
 reverse primer

<400> 29
 tacgccatcg ggcattgt 18

 <210> 30
 <211> 18
 <212> DNA
 <213> Burkholderia cepacia

 <220>
 <223> B. cepacia Genomovar I specific recA forward
 primer (BCRG11)

 <400> 30
 caggtcgtct ccacgggt 18

 <210> 31
 <211> 19
 <212> DNA
 <213> Burkholderia cepacia

 <220>
 <223> B. cepacia Genomovar I specific recA reverse
 primer (BCRG12)

 <400> 31
 cacgccgata ttcatacga 19

 <210> 32
 <211> 19
 <212> DNA
 <213> Burkholderia cepacia

 <220>
 <223> B. cepacia Genomovar III, RG-A specific recA
 forward primer (BCRG31)

 <400> 32
 gctcgacgtt caatatgcc 19

 <210> 33
 <211> 18
 <212> DNA
 <213> Burkholderia cepacia

 <220>
 <223> B. cepacia Genomovar III, RG-A specific recA
 reverse primer (BCRG32)

 <400> 33
 tcgagacgca ccgacgag 18

 <210> 34
 <211> 18

<212> DNA
 <213> Burkholderia cepacia

 <220>
 <223> B. cepacia recA reverse primer (BCR4)

 <400> 34
 gcgcagcgcc tgcgacat 18

 <210> 35
 <211> 18
 <212> DNA
 <213> Burkholderia cepacia

 <220>
 <223> B. cepacia recA forward primer (BCR3)

 <400> 35
 gtcgcaggcg ctgcgcaa 18

 <210> 36
 <211> 19
 <212> DNA
 <213> Burkholderia cepacia

 <220>
 <223> B. cepacia recA forward primer (BCRU1)

 <400> 36
 atcatgcgga tgggcgacg 19

 <210> 37
 <211> 19
 <212> DNA
 <213> Burkholderia cepacia

 <220>
 <223> B. cepacia recA reverse primer (BCRU2)

 <400> 37
 cagttctgtc gcttgatcg 19

 <210> 38
 <211> 19
 <212> DNA
 <213> Burkholderia cepacia

 <220>
 <223> B. cepacia Genomovar III, RG-B recA specific
 forward primer

 <400> 38
 gctgcaagtc atcgctgaa 19

<210> 39
<211> 18
<212> DNA
<213> Burkholderia cepacia

<220>
<223> B. cepacia RG-C specific recA forward primer

<400> 39
gtcgggtaaa accacgtg 18

<210> 40
<211> 18
<212> DNA
<213> Burkholderia cepacia

<220>
<223> B. cepacia RG-C specific recA reverse primer

<400> 40
tccgcagccg caccttca 18